



# PRODUCTION LICK

## 8% Urea + Phosphorus

Suitable for young growing cattle (150kg +) throughout the year when adequate feed is available.  
**This lick must be protected from water or rain as the product contains urea and is palatable.**

**Nutritional Analysis per Kg (as fed basis)**

ACTIVE	As fed %
Min. Crude Protein	10.5%
Min. Equivalent Protein	27.5%
Min. Total Protein	38%
Urea	8%
Molasses	1.5%
Calcium	8%
Phosphorus	6.5%
Sulphur	1.5%
Magnesium	0.8%
Salt	15%
Crude Fat (EE)	3.3%

**Added micro minerals:** Zinc, manganese, copper, iodine, cobalt and selenium. (organic & inorganic)

**Added vitamins & metabolite:** A, D3, E, Biotin (B7) and 25-hydroxycholecalciferol.

Contains lasalocid sodium 118.75 ppm/Kg

**Daily Intakes per head per day:**

Animal class - Cattle	Grams/hd/day
Grower 150Kg+	60-100
Grower 250+	60-150
Dry stock and bulls	150-250
Lactating cows	250-400

**Feeding Directions:** Ensure that animals are not starving when introduced to this product. Allow continuous access once feeding has commenced. Provide clean, fresh water at all times.

**Safety Directions:** Avoid contact with eyes and skin. Avoid breathing dust.

**First Aid:** Call Poisons Information Centre for advice 24 hrs (13 11 26) or nearest doctor at once. If contact with eyes occurs, wash out immediately with clean water. If contact

with skin or hair occurs, remove contaminated clothing and flush skin with running water.

**DO NOT FEED TO HORSES AND OTHER MONOGASTRICS. THIS PRODUCT MAY BE FATAL.**

**Keep out of reach of children. For animal treatment only.**

**With holding Period: Meat = Nil, Milk = Nil. Store below 30°C (Room Temp) in a dry place. Best used within 6 months.**

**Warning: THIS PRODUCT CONTAINS UREA – UREA CAN BE TOXIC TO LIVESTOCK & OTHER ANIMALS**

\*Products containing urea and other non-protein-nitrogen can be toxic to livestock causing illness and or death. Do not feed to starving, hungry or emaciated cattle and ensure that adequate fibre is always available. Livestock management and environmental conditions vary between locations and consideration of these dangers must be managed correctly. Urea supplementation should commence before animals lose weight, but effective urea supplementation can reduce rate of weight loss and promote feed efficiency, promoting animal weight gain. Effective supplementation can increase pasture or dry matter intake.

\*The success of this product/supplement is dependent on other production and nutritional factors being managed correctly. Most important is the supply of energy and protein through quality feed as well as good animal husbandry.

\*The use of this product after distribution or sale is beyond the control of **Morgan Feed Supplies**. Therefore, no warranty is given nor shall be implied for its suitability for any purpose. The manufacturer and their employees/contractors accept no responsibility for any consequence arising from the above-mentioned guides & product.

**Medicated animal food product**  
**This product does not contain restricted animal material**